

missed MIs represent a major cause of legal liability. Similarly, the advice to interpret an elevated troponin level in the clinical context is wise, yet may have a limited effect—estimates of pre-test probability have a high interobserver variability (3). Physicians often do not make formal quantitative calculations in practice (4) and make errors when they do (5).

Troponins, as currently used, have an excellent negative predictive value, but poor positive predictive value. The absolute value of troponin and the degree of increase on sequential testing are helpful in diagnosing acute MI, but may be less helpful in confirming thrombotic ACS as the cause. We have observed cases of acute troponin I increases as great as 10 ng/ml with stroke, diabetic ketoacidosis, and heart failure. As the key therapeutic decision to provide antithrombotics or urgent revascularization usually occurs on presentation, and such therapies can only benefit patients with significant ACS, practicing clinicians need a second confirmatory test for ACS. Such a test should be less severely affected by sepsis, tachyarrhythmia, hypotension, renal failure, chronic coronary disease, demand ischemia, heart failure, or cardiomyopathy, even if it is insensitive to small amounts of myonecrosis. That test, arguably, is the creatine kinase–myocardial band and index.

***Arnold Seto, MD, MPA**
David Tehrani, BS

*University of California
Irvine Medical Center
101 The City Drive
Orange, California 92868
E-mail: arnoldseto@yahoo.com

<http://dx.doi.org/10.1016/j.jacc.2013.01.045>

REFERENCES

1. Newby LK, Jesse RL, Babb JD, et al. ACCF 2012 Expert consensus document on practical clinical considerations in the interpretation of troponin elevations: a report of the American College of Cardiology Foundation Task Force on Clinical Expert Consensus Documents. *J Am Coll Cardiol* 2012;60:2427–63.
2. Waxman DA, Hecht S, Schappert J, Husk G. A model for troponin I as a quantitative predictor of in-hospital mortality. *J Am Coll Cardiol* 2006;48:1755–62.
3. Panju AA, Hemmelgarn BR, Guyatt GH, Simel DL. The rational clinical examination. Is this patient having a myocardial infarction? *JAMA* 1998;280:1256–63.
4. Reid MC, Lane DA, Feinstein AR. Academic calculations versus clinical judgments: practicing physicians' use of quantitative measures of test accuracy. *Am J Med* 1998;104:374–80.
5. Steurer J, Fischer JE, Bachmann LM, Koller M, ter Riet G. Communicating accuracy of tests to general practitioners: a controlled study. *BMJ* 2002;324:824–6.

Reply

The American College of Cardiology Foundation (ACCF) welcomes letters to inform its ongoing work and encourages such correspondence about its clinical policy statements. Because the ACCF document development process is rigorous and involves several layers of review by the writing committee, external peer reviewers, and participating organizations in the document, it cannot respond to each issue raised after a document has been published. The information, however, is forwarded to the Writing Committee co-chairs and oversight Task Force for review. If any issues are deemed by the College to affect patient safety, they will be considered immediately. Otherwise, the information will be considered during the next revision of the document.

***The American College of Cardiology Foundation**

*American College of Cardiology
2400 N Street, NW
Washington, DC 20037

<http://dx.doi.org/10.1016/j.jacc.2013.01.028>